Tutte polynomial of a fractal scale-free lattice *

Hanlin Chen,Yuanhua Liao, Hanyuan Deng†

College of Mathematics and Computer Science,

Hunan Normal University, Changsha, Hunan 410081, P. R. China

arXiv:1410.7988v1 [math.CO] 29 Oct 2014

Abstract

The Tutte polynomial of a graph, or equivalently the q-state Potts model partition function, is a two-variable polynomial graph invariant of considerable importance in both combinatorics and statistical physics. The computation of this invariant for a graph is NP-hard in general. In this paper, based on their self-similar structures, we recursively describe the Tutte polynomials of an infinite family of scale-free lattices. Furthermore, we give some exact analytical expressions of the Tutte polynomial for several special points at (X, Y)-plane.

Keywords: Potts model; Tutte polynomial; Spanning tree; Asymptotic growth constant; Self-similar.

1 Introduction

The Tutte polynomial $T(G; x, y)$ of a graph G, due to W. T. Tutte [1], is a two-variable polynomial and is deeply connected with many areas of both physics and mathematics. It is defined as

$$T(G; x, y) = \sum_{H \subseteq G} (x-1)^{r(G)-r(H)} (y-1)^{n(H)},$$

where the sum runs over all the spanning subgraphs H of G, $r(G) = |V(G)| - k(G)$, $n(G) = |E(G)| - |V(G)| + k(G)$ and $k(G)$ is the number of components of G. For a thorough survey on the Tutte polynomial, we refer the reader to [2, 3, 4]. The importance of the Tutte polynomial comes from the rich information it contains about the underlying graph. It contains several other polynomial invariants, such as the chromatic polynomial, the flow polynomial, the Jones polynomial and the all terminal reliability polynomial as partial evaluations, and various numerical invariants such as the number of spanning trees as complete evaluations. Furthermore, the Tutte polynomial has been widely studied in the field of statistical physics where it appears as the partition function of the zero-field

*Project supported by Hunan Provincial Natural Science Foundation of China (13JJ3053).

†Corresponding author: hydeng@hunnu.edu.cn

Potts model [5, 6]. In fact, let G be a graph with n vertices and $k(G)$ components and let Z_{zero} denote the partition function of the zero-field Potts model, then $Z_{zero}(G; q, v)$ and $T(G; x, y)$ satisfy the following relation [7, 8]:

$$Z_{zero}(G; q, v) = q^{k(G)} v^{n-k(G)} T(G; (q+v)/v, v+1).$$

In both fields of combinatorics and statistical physics, Tutte polynomials of many graphs (or lattices) have been computed by different methods [9, 10, 11, 12, 13, 14, 15, 16, 17]. Recently, on the basis of the subgraph expansion definition of the Tutte polynomial, a very useful method for computing the Tutte polynomial, called the subgraph-decomposition method, was developed by Donno *et al.* [18]. This technique is highly suited for computing the Tutte polynomial of self-similar graphs, and some applications of it can be found in [19, 20, 21].

The lattice (network) under consideration was introduced by Kaufman *et al.* [22, 23] and was further studied by Zhang *et al.* [24, 25] from the viewpoint of complex networks. The aim of this paper is to compute the Tutte polynomial of this deterministic scale-free network. For this purpose, we first partition the set of spanning subgraphs of this network into two disjoint subsets. In this way, we can express the Tutte polynomial by two summands. Then, we study the contributions of all kinds of possible combinations of this two subsets. Finally, base on the self-similar structure of the network, we obtain a recursive formula for each summand. In particular, as special cases of the general Tutte polynomial, we get:

- the recursive formula for computing the Tutte polynomial $T(G; x, y)$;
- the number $\tau(G)$ of spanning trees and the asymptotic growth constant $\lim\limits_{n\to\infty} \frac{ln\tau(G_n)}{|v(G_n)|}$;
- the dimension of the bicycle space;
- the number of acyclic root-connected orientations; the number of indegree sequences of strongly connected orientations.

2 Preliminaries

In this section, we briefly discuss some necessary background that will be used for our calculations. We use standard graph terminology and the words "network" and "graph"

indistinctly. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The vertices a and b are the end-points of an edge $\{a, b\}$. An *orientation* of graph G is the digraph defined by the choice of a direction for every edge of $E(G)$. A *directed cycle* of a digraph is a set of edges forming a cycle of the graph such that they are all directed accordingly with a direction for the cycle. A digraph is *acyclic* if it has no directed cycle, and *strongly connected* if for every pair of vertices there is a directed cycle containing them. A *sink* for a digraph is a vertex with no outgoing edge.



Figure 1: The lattices G_0, G_1, G_2 and G_3.

The network, as shown in Figure 1, can be constructed as follows.

For $n = 0$, G_0 is the complete graph K_2. For $n \geq 0$, G_{n+1} can be constructed from four copies of G_n by merging four groups of vertices and adding a new edge. Specifically, let X_n and Y_n, hereafter called special vertices of G_n, be the leftmost and the rightmost vertex of G_n. X_n and X_n are combined into the special vertex X_{n+1} of G_{n+1}, Y_n and Y_n are combined into the special vertex Y_{n+1} of G_{n+1}, and a new edge e_n is added between two vertices combined by Y_n and X_n. The construction of G_{n+1} is illustrated in Figure 2.



Figure 2: The construction of G_{n+1}

3 Recursive formulas

In this section, we give the computational formulas of Tutte polynomial of the lattice G_n. It is easy to obtain that the order and size of the lattice G_n are, respectively,

$$|V(G_n)| = \frac{2 \times 4^n + 4}{3}, \quad |E(G_n)| = \frac{4^{n+1} - 1}{3}.$$

And the average degree after n iterations is $\langle \mathrm{k} \rangle_n = \frac{2|E(G_n)|}{|V(G_n)|}$, which approaches 4 in the infinite n limit.

To investigate the Tutte polynomial $T(G_n; x, y)$. First of all, we partition the set of the spanning subgraph of G_n into two disjoint subsets:

- $\mathcal{G}_{1,n}$ denotes the set of spanning subgraphs of G_n, where two special vertices X_n and Y_n of G_n belong to the same component;
- $\mathcal{G}_{2,n}$ denotes the set of spanning subgraphs of G_n, where two special vertices X_n and Y_n of G_n do not belong to the same component.



Figure 3: The spanning subgraph of G_n: Type I (left), Type II (right).

Observe that, for each $n \geq 0$, we have the partition $\mathcal{G}_{1,n} \cup \mathcal{G}_{2,n}$ of the set of spanning subgraphs of G_n. Next, let $T_n(x, y)$ denote the Tutte polynomial $T(G_n; x, y)$ of G_n. For every $n \geq 1$, $T_{1,n}(x, y)$ and $T_{2,n}(x, y)$ are the following polynomials:

- $T_{1,n}(x, y) = \sum_{H \in \mathcal{G}_{1,n}} (x-1)^{r(G_n)-r(H)}(y-1)^{n(H)}$;
- $T_{2,n}(x, y) = \sum_{H \in \mathcal{G}_{2,n}} (x-1)^{r(G_n)-r(H)}(y-1)^{n(H)}$.

We have

$$T_n(x, y) = T_{1,n}(x, y) + T_{2,n}(x, y).$$

In order to obtain $T_n(x, y)$, we need to find recursive formulas on $T_{1,n}(x, y)$ and $T_{2,n}(x, y)$. For this purpose, we analyze the relation between spanning subgraphs of G_{n+1} and spanning subgraph of G_n. Observe from Figure 2 that G_{n+1} is constructed from four copies of G_n by merging some special vertices and adding a new edge e_n. Thus, a spanning subgraphs of G_{n+1} is combined by the spanning subgraphs of the four copies

$G_n^i (i = 1, 2, 3, 4)$ of G_n with S, where S may be $\{e_n\}$ or $\emptyset$. Indeed, a spanning subgraph H of G_{n+1} is uniquely determined by the restriction of H to the four copies G_n^i (denoted by $H_i (i = 1, 2, 3, 4)$, respectively) and S, and vice versa. Therefore, the Tutte polynomial of G_{n+1} can be written as

$$T_{n+1}(x, y) = \sum_{H_i \subseteq G_n^i; (\bigcup_{i=1}^{4} H_i) \bigcup S = H} (x-1)^{r(G_{n+1})-r(H)}(y-1)^{n(H)}.$$

where the sum runs over all spanning subgraphs H_i of G_n^i $(i = 1, 2, 3, 4)$ and S. Now, we need to know how $r(H)$ and $n(H)$ depend on $r(H_i)$ and $n(H_i)$, for $i = 1, 2, 3, 4$. Note that $|V(G_{n+1})| = 4|V(G_n)| - 4$, $|E(H)| = \sum_{i=1}^{4} |E(H_i)| + 1$ if $S = \{e_n\}$ and $|E(H)| = \sum_{i=1}^{4} |E(H_i)|$ if $S = \emptyset$. So, there are two cases to be considered.

Case 1. $S = \{e_n\}$.

In this case, we consider the spanning subgraph H of G_{n+1}, which contains the new adding edge $\{e_n\}$, and $|E(H)| = \sum_{i=1}^{4} |E(H_i)| + 1$.

Subcase 1. If $k(H) = \sum_{i=1}^{4} k(H_i) - 3$, then

$$r(H) = |V(H)| - k(H) = (4|V(G_n)| - 4) - (\sum_{i=1}^{4} k(H_i) - 3) = \sum_{i=1}^{4} r(H_i) - 1.$$

Moreover, we have

$$n(H) = |E(H)| - r(H) = (\sum_{i=1}^{4} |E(H_i)| + 1) - (\sum_{i=1}^{4} r(H_i) - 1) = \sum_{i=1}^{4} n(H_i) + 2$$

and

$$r(G_{n+1}) - r(H) = (|V(G_{n+1})| - 1) - (\sum_{i=1}^{4} r(H_i) - 1) = \sum_{i=1}^{4} (r(G_n) - r(H_i)).$$

Thus,

$$(x-1)^{r(G_{n+1})-r(H)}(y-1)^{n(H)} = (y-1)^2 \prod_{i=1}^{4} (x-1)^{r(G_n)-r(H_i)}(y-1)^{n(H_i)}.$$

Subcase 2. If $k(H) = \sum_{i=1}^{4} k(H_i) - 4$, then

$$r(H) = |V(H)| - k(H) = (4|V(G_n)| - 4) - (\sum_{i=1}^{4} k(H_i) - 4) = \sum_{i=1}^{4} r(H_i).$$

Moreover, we have

$$n(H) = |E(H)| - r(H) = (\sum_{i=1}^{4} |E(H_i)| + 1) - (\sum_{i=1}^{4} r(H_i)) = \sum_{i=1}^{4} n(H_i) + 1$$

and

$$r(G_{n+1}) - r(H) = (|V(G_{n+1})| - 1) - (\sum_{i=1}^{4} r(H_i)) = \sum_{i=1}^{4}(r(G_n) - r(H_i)) - 1.$$

Hence,

$$(x-1)^{r(G_{n+1})-r(H)}(y-1)^{n(H)} = \frac{y-1}{x-1}\prod_{i=1}^{4}(x-1)^{r(G_n)-r(H_i)}(y-1)^{n(H_i)}.$$

Subcase 3. If $k(H) = \sum\limits_{i=1}^{4} k(H_i) - 5$, then we can obtained, similarly, that

$$(x-1)^{r(G_{n+1})-r(H)}(y-1)^{n(H)} = \frac{1}{(x-1)^2}\prod_{i=1}^{4}(x-1)^{r(G_n)-r(H_i)}(y-1)^{n(H_i)}.$$

Case 2. $S = \emptyset$.

In this case, we consider the spanning subgraph H of G_{n+1}, which does not contain the new adding edge e_n, and $|E(H)| = \sum\limits_{i=1}^{4} |E(H_i)|$.

Subcase 1. If $k(H) = \sum\limits_{i=1}^{4} k(H_i) - 3$, then

$$r(H) = |V(H)| - k(H) = (4|V(G_n)| - 4) - (\sum_{i=1}^{4} k(H_i) - 3) = \sum_{i=1}^{4} r(H_i) - 1.$$

Moreover, we have

$$n(H) = |E(H)| - r(H) = (\sum_{i=1}^{4} |E(H_i)|) - (\sum_{i=1}^{4} r(H_i) - 1) = \sum_{i=1}^{4} n(H_i) + 1$$

and

$$r(G_{n+1}) - r(H) = (|V(G_{n+1})| - 1) - (\sum_{i=1}^{4} r(H_i) - 1) = \sum_{i=1}^{4}(r(G_n) - r(H_i))$$

Thus,

$$(x-1)^{r(G_{n+1})-r(H)}(y-1)^{n(H)} = (y-1)\prod_{i=1}^{4}(x-1)^{r(G_n)-r(H_i)}(y-1)^{n(H_i)}.$$

Subcase 2. If $k(H) = \sum\limits_{i=1}^{4} k(H_i) - 4$, then

$$r(H) = |V(H)| - k(H) = (4|V(G_n)| - 4) - (\sum_{i=1}^{4} k(H_i) - 4) = \sum_{i=1}^{4} r(H_i).$$

Moreover, we have

$$n(H) = |E(H)| - r(H) = (\sum_{i=1}^{4} |E(H_i)|) - (\sum_{i=1}^{4} r(H_i)) = \sum_{i=1}^{4} n(H_i)$$

and

$$r(G_{n+1}) - r(H) = (|V(G_{n+1})| - 1) - (\sum_{i=1}^{4} r(H_i)) = \sum_{i=1}^{4} (r(G_n) - r(H_i)) - 1$$

Thus we have

$$(x-1)^{r(G_{n+1})-r(H)}(y-1)^{n(H)} = \frac{1}{x-1}\prod_{i=1}^{4}(x-1)^{r(G_n)-r(H_i)}(y-1)^{n(H_i)}.$$

For the convenience of discussion, we use solid lines to join two special vertices when the corresponding spanning subgraph of G_n belongs to $\mathcal{G}_{1,n}$; Otherwise, we use dotted lines instead of solid lines. We distinguish two types of the spanning subgraph as shown in Figure 3.

Theorem 1. *The Tutte polynomial $T_{n+1}(x,y)$ of G_{n+1} is given by*

$$T_{n+1}(x,y) = T_{1,n+1}(x,y) + T_{2,n+1}(x,y),$$

where the polynomials $T_{1,n+1}(x,y)$ and $T_{2,n+1}(x,y)$ satisfy the following recursive relations:

$$T_{1,n+1}(x,y) = y(y-1)T_{1,n}^4 + \frac{4y}{x-1}T_{1,n}^3 T_{2,n} + \frac{2x+2}{(x-1)^2}T_{1,n}^2 T_{2,n}^2, \tag{1}$$

$$T_{2,n+1}(x,y) = \frac{2y+2}{x-1}T_{1,n}^2 T_{2,n}^2 + \frac{4x}{(x-1)^2}T_{1,n}T_{2,n}^3 + \frac{x}{(x-1)^2}T_{2,n}^4 \tag{2}$$

with initial conditions $T_{1,0}(x,y) = 1,\ T_{2,0}(x,y) = x-1$.

Proof. The initial conditions are easily verified. The strategy of the proof is to study all possible configurations of the spanning subgraph H_i in G_n^i $(i = 1,2,3,4)$, and analyze which kind of contributions they give to $T_{1,n}(x,y)$ and $T_{2,n}(x,y)$. As shown in Table 1, a configuration produces a basic term of form $T_{1,n}^i T_{2,n}^j (i+j=4)$, and by the previous analysis, each basic term has to be multiplied by a factor $(y-1)^2$, $\frac{y-1}{x-1}$, $\frac{1}{(y-1)^2}$ or $y-1$, $\frac{1}{x-1}$ according to Case 1 and Case 2. More precisely, (i) the factor is $(y-1)^2$ if $S = \{e_n\}$ and $k(H) = \sum_{i=1}^{4} k(H_i) - 3$; (ii) the factor is $\frac{y-1}{x-1}$ if $S = \{e_n\}$ and $k(H) = \sum_{i=1}^{4} k(H_i) - 4$; (iii) the factor is $\frac{1}{(x-1)^2}$ if $S = \{e_n\}$ and $k(H) = \sum_{i=1}^{4} k(H_i) - 5$; (iv) the factor is $(y-1)$ if $S = \emptyset$ and $k(H) = \sum_{i=1}^{4} k(H_i) - 3$; (v) the factor is $\frac{1}{x-1}$ if $S = \emptyset$ and $k(H) = \sum_{i=1}^{4} k(H_i) - 4$.

Table 1: All combinations and corresponding contributions.

Configuration	S	Contribution	Type	Configuration	S	Contribution	Type
	$\{e_n\}$	$(y-1)^2T_{1,n}^4$	I		$\emptyset$	$(y-1)T_{1,n}^4$	I
	$\{e_n\}$	$\frac{y-1}{x-1}T_{1,n}^3T_{2,n}$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^3T_{2,n}$	I
	$\{e_n\}$	$\frac{y-1}{x-1}T_{1,n}^3T_{2,n}$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^3T_{2,n}$	I
	$\{e_n\}$	$\frac{y-1}{x-1}T_{1,n}^3T_{2,n}$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^3T_{2,n}$	I
	$\{e_n\}$	$\frac{y-1}{x-1}T_{1,n}^3T_{2,n}$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^3T_{2,n}$	I
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}^2T_{2,n}^2$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^2T_{2,n}^2$	I
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}^2T_{2,n}^2$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^2T_{2,n}^2$	I
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}^2T_{2,n}^2$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^2T_{2,n}^2$	II
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}^2T_{2,n}^2$	I		$\emptyset$	$\frac{1}{x-1}T_{1,n}^2T_{2,n}^2$	II
	$\{e_n\}$	$\frac{y-1}{x-1}T_{1,n}^2T_{2,n}^2$	II		$\emptyset$	$\frac{1}{x-1}T_{1,n}^2T_{2,n}^2$	II
	$\{e_n\}$	$\frac{y-1}{x-1}T_{1,n}^2T_{2,n}^2$	II		$\emptyset$	$\frac{1}{x-1}T_{1,n}^2T_{2,n}^2$	II
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}T_{2,n}^3$	II		$\emptyset$	$\frac{1}{x-1}T_{1,n}T_{2,n}^3$	II
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}T_{2,n}^3$	II		$\emptyset$	$\frac{1}{x-1}T_{1,n}T_{2,n}^3$	II
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}T_{2,n}^3$	II		$\emptyset$	$\frac{1}{x-1}T_{1,n}T_{2,n}^3$	II
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{1,n}T_{2,n}^3$	II		$\emptyset$	$\frac{1}{x-1}T_{1,n}T_{2,n}^3$	II
	$\{e_n\}$	$\frac{1}{(x-1)^2}T_{2,n}^4$	II		$\emptyset$	$\frac{1}{x-1}T_{2,n}^4$	II

From Table 1, we can establish Eqs.(1)-(2), and the proof is completed. □

According to Eq.(2), it is easy to prove by induction that $x-1$ divides $T_{2,n}(x,y)$ in $\mathbb{Z}[x,y]$. Thus, we can rewrite $T_{2,n}(x,y)$ as $(x-1)N_n(x,y)$ in $\mathbb{Z}[x,y]$, and Theorem 1 can be reduced to the following:

Theorem 2. *The Tutte polynomial $T_{n+1}(x,y)$ of G_{n+1} is given by*

$$T_{n+1}(x,y) = T_{1,n+1}(x,y) + (x-1)N_{n+1}(x,y),$$

where the polynomial $T_{1,n+1}(x,y)$, $N_{n+1}(x,y)$ *satisfy the following recursive relations:*

$$T_{1,n+1}(x,y) = y(y-1)T_{1,n}^4 + 4yT_{1,n}^3N_n + (2x+2)T_{1,n}^2N_n^2,$$
$$N_{n+1}(x,y) = (2y+2)T_{1,n}^2N_n^2 + 4xT_{1,n}N_n^3 + x(x-1)N_n^4$$

with initial conditions $T_{1,0}(x,y) = 1,\ N_0(x,y) = 1.$

It is well-known that the evaluation of the Tutte polynomial for a particular for a particular point at (X,Y)-plane is related to some combinatorial information and algebraic properties of the graph considered.

(1) $T(G;1,0)$ = the number of acyclic root-connected trees of G;

(2) $T(G;1,1) = \tau(G)$, i.e., the number of spanning trees;

(3) $T(G;0,1)$ = the number of indegree sequences of strongly connected orientations of G;

(4) $T(G;-1,-1) = (-1)^{|E(G)|}(-2)^{dim(\mathcal{B})}$, where $\mathcal{B}$ is the bicyclic space of G.

Theorem 3. *The number of acyclic root-connected orientations of* G_n *is given by* $T_n(1,0) = \prod_{i=0}^{n}(i+1)^{2\times 4^{n-i}}$; *The number of indegree sequences of strongly connected orientations of* G_n *is given by* $T_n(0,1) = \frac{n}{2}\prod_{i=0}^{n}(i+1)^{2\times 4^{n-i}}$.

Proof. By taking $x=1$ and $y=0$ in Theorem 2, we have $T_n(1,0) = T_{1,n}(1,0)$, and

$$T_{1,n}(1,0) = 4T_{1,n-1}^2(1,0)N_n^2(1,0), \tag{3}$$

$$N_n(1,0) = 2T_{1,n-1}^2(1,0)N_{n-1}^2(1,0) + 4T_{1,n-1}(1,0)N_{n-1}^3(1,0) \tag{4}$$

A useful relation yields from Eqs.(3) and (4)

$$\frac{N_n(1,0)}{T_{1,n}(1,0)} = \frac{1}{2} + \frac{N_{n-1}(1,0)}{T_{1,n-1}(1,0)}. \tag{5}$$

It implies that

$$\frac{N_n(1,0)}{T_{1,n}(1,0)} = \frac{n}{2} + \frac{N_0(1,0)}{T_{1,0}(1,0)},$$

and

$$N_n(1,0) = \frac{n+2}{2}T_{1,n}(1,0) \tag{6}$$

since $T_0(1,0) = 1,\ N_0(1,0) = 1.$

Substituting (6) into (3) and using the initial condition $T_{1,0}(1,0)=1$, we have

$$T_{1,n}(1,0)=(n+1)^2T_{1,n-1}^4(1,0)=\prod_{i=0}^{n}(i+1)^{2\times 4^{n-i}}.$$

Similarly, by taking $x=0$ and $y=1$ in Theorem 2, we have $T_n(0,1)=T_{1,n}(0,1)-N_n(0,1)$, and

$$\begin{aligned} T_{1,n}(0,1) &= 4T_{1,n-1}^3(0,1)N_n(0,1)+2T_{1,n-1}^2(0,1)N_{n-1}^2(0,1),\\ N_n(0,1) &= 4T_{1,n-1}^2(0,1)N_{n-1}^2(0,1). \end{aligned}$$

Using the same techniques, we can obtained

$$T_{1,n}(0,1)=\frac{n+2}{2}N_n(0,1),\ \ N_n(0,1)=\prod_{i=0}^{n}(i+1)^{2\times 4^{n-i}}.$$

And

$$T_n(0,1)=\frac{n+2}{2}N_n(0,1)-N_n(0,1)=\frac{n}{2}N_n(0,1)=\frac{n}{2}\prod_{i=0}^{n}(i+1)^{2\times 4^{n-i}}.$$

□

From above, the number $T_n(0,1)$ of acyclic root-connected trees and the number $T_n(1,0)$ of indegree sequences of strongly connected orientations in G_n are satisfied $T_n(0,1)=\frac{n}{2}T_n(1,0)$.

Theorem 4. *For a positive integer $n\geqslant 1$, the Tutte polynomial of G_n, $T_n(x,y)$ along the line $y=x$ is given by*

$$T_n(x,x)=x(x^2+5x+2)^{\frac{4^n-1}{3}}.$$

Proof. By taking $y=x$ in Theorem 2, we have

$$T_{1,n}(x,x)=x(x-1)T_{1,n-1}^4+4xT_{1,n-1}^3N_{n-1}+(2x+2)T_{1,n-1}^2N_{n-1}^2,$$

$$N_n(x,x)=(2x+2)T_{1,n-1}^2N_{n-1}^2+4xT_{1,n-1}N_{n-1}^3+x(x-1)N_{n-1}^4, \tag{7}$$

and $T_{1,0}(x,x)=N_0(x,x)=1$. It can be obtained easily that $T_{1,n}(x,x)=N_n(x,x)$ by induction. Substituting it into Eq.(7) and using the initial condition $N_0(x,x)=1$, we have

$$N_n(x,x)=(x^2+5x+2)N_{n-1}^4=(x^2+5x+2)^{\frac{4^n-1}{3}}.$$

Thus, $T_n(x,x) = T_{1,n}(x,x) + (x-1)N_n(x,x) = xN_n(x,x) = x(x^2+5x+2)^{\frac{4^n-1}{3}}$. □

Since the number of spanning trees is $\tau(G) = T(G;1,1)$, from Theorem 4, we can obtain immediately the following result, which was also obtained in [25] by employing the decimation teachnique.

Corollary 5. *The number of spanning trees of G_n is given by $T_n(1,1) = 8^{\frac{4^n-1}{3}} = 2^{4^n-1}$. The asymptotic growth constant is*

$$\lim_{n\to\infty} \frac{ln\tau(G_n)}{|V(G_n)|} = \frac{3}{2} ln2 \simeq 1.0397.$$

Similarly, $T_n(-1,-1) = (-1)\times(-2)^{\frac{4^n-1}{3}} = (-1)^{\frac{4^{n+1}-1}{3}}(-2)^{\frac{4^n-1}{3}} = (-1)^{|E(G_n)|}(-2)^{dim(\mathcal{B})}$ by taking $x=-1$ in Theorem 4. So, we have

Corollary 6. *The dimension of the bicycle space of G_n is $\frac{4^n-1}{3}$.*

Appendices

4 Tutte polynomial of two other scale-free networks

In this section, we consider the Tutte polynomial of two scale-free networks (2,2)-flower and (1,3)-flower which have some similar characteristics: identical degree sequences, without crossing edges and always connected. Most of the topological properties of (2,2)-flower and (1,3)-flower can be determined exactly [26].

4.1 (2,2)-flower

If the newly added edge is ignored at each iterative generation, then the fractal lattice G_n considered above become the (2,2)-flower F_n, see Figure 4. And the contributions to $T_{1,n}(x,y)$ and $T_{2,n}(x,y)$ are degraded into the case of $S=\emptyset$, and listed on the right of Table 1. So, we can obtain

$$T_n(x,y) = T_{1,n}(x,y) + (x-1)N_n(x,y),$$

where the polynomials $T_{1,n}(x,y)$, $N_n(x,y)$ satisfy the following recursive relations:

$$\begin{aligned} T_{1,n}(x,y) &= (y-1)T_{1,n-1}^4 + 4T_{1,n-1}^3 N_{n-1} + 2(x-1)T_{1,n-1}^2 N_{n-1}^2, \\ N_n(x,y) &= 4T_{1,n-1}^2 N_{n-1}^2 + 4(x-1)T_{1,n-1}N_{n-1}^3 + (x-1)^2 N_{n-1}^4. \end{aligned}$$



Figure 4: Construction method of the (2,2)-flower F_n.

If $x = y = 1$, then $T_n(1,1) = T_{1,n}(1,1)$ and $T_{1,n}(1,1) = N_n(1,1)$. Thus, $T_n(1,1) = 4T^4_{n-1}(1,1)$. Since the initial value $T_0(1,1) = 1$, we can obtain

$$\tau(F_n) = T_n(1,1) = 2^{\frac{2}{3}(4^n-1)}$$

and

$$\lim_{n\to\infty} \frac{ln\tau(F_n)}{|V(F_n)|} = ln2 \simeq 0.6931.$$

4.2 (1,3)-flower



Figure 5: Construction method of the (1,3)-flower F_n.

For the (1,3)-flower F_n, see Figure 5, choosing two adjacent vertices with largest degree as special vertices in each iteration, we can obtain similarly the following recursive relation:

$$T_n(x,y) = T_{1,n}(x,y) + T_{2,n}(x,y),$$

where the polynomials $T_{1,n}(x,y)$ and $T_{2,n}(x,y)$ satisfy the following recursive relations:

$$\begin{aligned} T_{1,n}(x,y) &= (y-1)T^4_{1,n-1} + \frac{4}{x-1}T^3_{1,n-1}T_{2,n-1} + \frac{3}{x-1}T^2_{1,n-1}T^2_{2,n-1} + \frac{1}{x-1}T_{1,n-1}T^3_{2,n-1}, \\ T_{2,n}(x,y) &= \frac{3}{x-1}T^2_{1,n-1}T^2_{2,n-1} + \frac{3}{x-1}T_{1,n-1}T^3_{2,n-1} + \frac{1}{x-1}T^4_{2,n-1}. \end{aligned}$$

with the initial conditions $T_{1,0}(x,y) = 1$, $T_{2,0}(x,y) = x-1$. And

$$T_n(x,y) = T_{1,n}(x,y) + (x-1)N_n(x,y),$$

where the polynomials $T_{1,n}(x,y)$ and $N_n(x,y)$ satisfy the following recursive relations:

$$\begin{aligned} T_{1,n}(x,y) &= (y-1)T_{1,n-1}^4 + 4T_{1,n-1}^3 N_{n-1} + 3(x-1)T_{1,n-1}^2 N_{n-1}^2 + (x-1)^2 T_{1,n-1} N_{n-1}^3, \\ N_n(x,y) &= 3T_{1,n-1}^2 N_{n-1}^2 + 3(x-1)T_{1,n-1} N_{n-1}^3 + (x-1)^2 N_{n-1}^4 \end{aligned}$$

with the initial conditions $T_{1,0}(x,y) = 1$ and $N_0(x,y) = 1$.

Similarly, if $x = y = 1$, then $T_n(1,1) = T_{1,n}(1,1)$ and

$$T_{1,n}(1,1) = 4T_{1,n-1}^3(1,1)N_{n-1}(1,1), \tag{8}$$

$$N_n(1,1) = 3T_{1,n-1}^2(1,1)N_{n-1}^2(1,1). \tag{9}$$

From Eqs.(8) and (9), we have

$$T_{1,n}(1,1) = (\frac{4}{3})^n N_n(1,1).$$

Since the initial value $N_0(1,1) = 1$, we can obtain

$$\tau(F_n) = T_n(1,1) = 3^{(4^n-3n-1)/9} 4^{(2\times 4^n+3n-2)/9}$$

and

$$\lim_{n\to\infty} \frac{ln\tau(F_n)}{|V(F_n)|} = \frac{1}{6}(4ln2 + ln3) \simeq 0.6452.$$

which are coincides with the results in [27] based on the relationship between determinants of submatrices of the Laplacian matrix.

References

[1] Tutte W T, 1954 *Canad. J. Math.* **6** 80

[2] Jaeger F, Vertigan D L and Welsh D J A, 1990 *Math. Proc. Cambridge Philos. Soc.* **108** 35

[3] Welsh D J A and Merino C, 2000 *J. Math. Phys.* **41** 1127

[4] Ellis-Monahan J and Merino C, 2011 Graph polynomials and their applications I: the Tutte polynomial *Structural Analysis of Complex Networks* ed D Matthias (Boston: Birkhüser)

[5] Potts R B, 1952 *Proc. Cambridge Philos. Soc.* **48** 106

[6] Wu F Y, 1992 *Rev. Mod. Phys.* **64** 1099

[7] Fortuin C M and Kasteleyn P W, 1972 *Physica* **57** 536

[8] Beaudin L, Ellis-Monahan J, Pangborn G and Shrock R, 2010 *Discrete Math.* **310** 2037

[9] Shrock R, 2000 *Physica A* **283** 388

[10] Chang S C and Shrock R, 2000 *Phys. A* **286** 189

[11] Chang S C and Shrock R, 2001 *Phys. A* **296** 234

[12] Chang S C and Shrock R, 2004 *Adv. Appl. Math.* **32** 44

[13] Chang S C and Shrock R, 2008 *J. Stat. Phys.* **130** 1011

[14] Shrock R, 2011 *J. Phys. A: Math. Theor.* **44** 145002

[15] Shrock R and Xu Y, 2012 *J. Phys. A: Math. Theor.* **45** 055212

[16] Salas J and Sokal A D, 2009 *J. Stat. Phys.* **135** 279

[17] Alvarez P D, Canfora F, Reyes S A and Riquelme S, 2012 *Eur. Phys. J. B* **85** 99

[18] Donno A and Iacono D, 2013 *Adv. Geom.* **13** 663

[19] Liao Y H, Hou Y P and Shen X L, 2013 *EPL* **104** 38001

[20] Liao Y H, Fang A X and Hou Y P, 2013 *Phys. A* **392** 4584

[21] Gong H L and Jin X A, 2014 *Phys. A* **414** 143

[22] Kaufman M and Griffiths R B, 1981 *Phys. Rev. B* **24** 496

[23] Griffiths R B and Kaufman M, 1982 *Phys. Rev. B* **26** 5022

[24] Zhang Z Z, Zhou S G, Zou T and Chen G S, 2008 *J. Stat. Mech.* P09008

[25] Zhang Z Z, Liu H X, Wu B and Zhou T, 2011 *Phys. Rev. E* **83** 016116

[26] Zhang Z Z, Zhou S G, Zou T, Chen L C and Guan J H, 2009 *Phys. Rev. E* **79** 031110

[27] Lin Y, Wu B, Zhang Z Z and Chen G, 2011 *J. Math. Phys.* **52** 113303